

**"an effective antirheumatic agent"\***

*nonhormonal anti-arthritic*

**BUTAZOLIDIN®**


(brand of phenylbutazone)

**relieves pain • improves function • resolves inflammation**

The standing of BUTAZOLIDIN among today's anti-arthritics is attested by more than 250 published reports. From this combined experience it is evident that BUTAZOLIDIN has achieved recognition as a potent agent capable of producing clinical results that compare favorably with those of the hormones.

Indications: Gouty Arthritis Rheumatoid Arthritis Psoriatic Arthritis  
Rheumatoid Spondylitis Painful Shoulder Syndrome  
BUTAZOLIDIN® (brand of phenylbutazone) red coated tablets of 100 mg.

\*Bunick, J. J.: Research Activities in Rheumatic Diseases, Pub. Health Rep. 69:437, 1954.



**GEIGY PHARMACEUTICALS**

Division of Geigy Chemical Corporation, 220 Church Street, New York 13, N.Y.

48555

Now . . . a totally new nonbarbiturate hypnotic-sedative!



for a good night's sleep with clear awakening

**DORIDEN<sup>®</sup>**

(glutethimide, CIBA)

In most cases—

Rapid onset—15-20 minutes

Lasts 4-8 hours

No hangover

Dosage: 0.25 to 0.5 Gm. before bedtime.

Scored 0.25- and 0.5-Gm. tablets.

CIBA, Summit, N.J.

2/2076M

- ✿ highly soluble in both acid and alkaline body fluids, especially at pH of renal tubules
- ✿ high plasma levels rapidly attained
- ✿ no alkalization or forcing of fluids required
- ✿ single sulfonamide - not a mixture
- ✿ no danger of secondary fungus infections
- ✿ wide antibacterial range

**you prescribe  
these distinctive features  
when you prescribe . . .**




**Hoffmann - La Roche Inc • Roche Park • Nutley 10 • N.J.**

**GANTRISIN®**—brand of sulfoxazole (3,4-dimethyl-5-sulfanilamido-isoxazole)

# CAMBRIDGE EDUCATIONAL CARDIOSCOPE

Bulletin 184



Overall Dimensions: 74" x 25" x 25"


**CAMBRIDGE INSTRUMENT CO., INC.**

374 Grand Central Terminal, New York, N. Y.  
PIONEER MANUFACTURERS OF THE ELECTROCARDIOGRAPH

## Send for BULLETIN 184

THE CAMBRIDGE EDUCATIONAL CARDIOSCOPE is used in many prominent Medical Schools for teaching cardiology by audio-visual demonstration. The electrocardiogram, heart sounds, vectorcardiogram, EKG, BCG, intra-cardiac pressures, etc., may be individually shown on a 16" cathode ray tube having a long persistence screen. While viewing the physiological data, the related heart sounds may be heard by an entire group through individual stethophones. *With the Cambridge Educational Cardioscope, true audio-visual teaching of Cardiology is now a reality—a must for the modern teaching program.*

the last tablet as accurate as the first



# sealed-in-foil CLINITEST® BRAND REAGENT TABLETS

a rapid, reliable urine-sugar test every time because every batch of *Clinitest* Sealed-in-Foil Reagent Tablets is tested for stability under conditions as exacting as a tropical rainy season—86° to 90° temperatures and 95% humidity.

*Clinitest* Reagent Tablets, Sealed in Foil, boxes of 24 and 500.

**AMES DIAGNOSTICS**  
Adjuncts in Clinical Management




**AMES COMPANY, INC. • ELKHART, INDIANA**  
Ames Company of Canada, Ltd., Toronto

# Meratran

(pipradrol hydrochloride)

a unique central motivant with demonstrated subcortical activity subtly returns your emotionally fatigued and depressed patients to their usual level of alertness, interest and productivity<sup>1,2</sup>.



Meratran's action, in easily adjusted doses, is prompt—subtle—comfortable—Its effectiveness is prolonged.


• no appreciable effect on blood pressure and respiration • restores needed sense of well-being • no tolerance or drug habituation • normal appetite undisturbed • no jitters - no apprehension • little or no insomnia • wide range of safety • no rebound letdown

## Rx INFORMATION

# Meratran

**Indications:** Emotional fatigue, unhappiness of more common type (financial worry, social stress). Situational stress or *mild* depression. Adjunctive therapy in certain psychoses and psycho-neuroses.<sup>3,4</sup>

**Composition:** alpha-(2-piperidyl) benzhydrol hydrochloride with the following structure:



**Dosage:** For emotional fatigue and mild depression, 1 to 6 mg. daily. Individual patient response must be observed and daily dosage and duration of administration adjusted to patient response.

**Supplied:** Small pink tablets containing: 1 mg. Meratran (pipradrol) hydrochloride.\* Bottles of 100.

\* Alpha (2 piperidyl) benzhydrol hydrochloride.

1. Heath, R. C.: Discussion of paper by Howard D. Fabing (alpha-(2-piperidyl) benzhydrol hydrochloride, a new central stimulant in the treatment of blepharospasm, spasmodic torticollis and narcolepsy) presented before the American Neurological Association, Atlantic City, June 14, 1954. 2. Fabing, H. D., Hawkins, J. R., Moulton, J. A. L.: Clinical studies on alpha-(2-piperidyl) benzhydrol hydrochloride, a new antidepressant drug. Presented before the American Psychiatric Association, St. Louis, Missouri, May 3, 1954. 3. Fabing, H. D.: Alpha-(2-piperidyl) benzhydrol hydrochloride, a new central stimulant, in the treatment of blepharospasm, spasmodic torticollis and narcolepsy, preliminary report (Motion Picture Demonstration), American Neurological Association, June 14, 1954. 4. Himwich, H. E.: Discussion of paper (reference 2), American Psychiatric Association, May 3, 1954.



**THE WM. S. MERRELL COMPANY**

New York • CINCINNATI • St. Thomas, Ontario  
PIONEER IN MEDICINE FOR OVER 125 YEARS

Thank You

We thank you, doctor, for prescribing Trevidal. Most of you who have employed this new, balanced protective antacid have continued to use it routinely. For those of you who have not yet had a chance to evaluate the unique antacid activity of Trevidal, may we suggest you write us today for a clinical trial supply.

## TREVIDAL®

### EACH TABLET CONTAINS:

Unique vegetable mucin  
supplies protective coat to  
irritated stomach lining



Regonol\*\* . . . . . 100 mg.

Balance of ingredients  
avoids constipation,  
diarrhea, or alkalosis



Magnesium trisilicate . . . . . 150 mg.  
Aluminum hydroxide gel . . . . . 90 mg.  
Calcium carbonate . . . . . 105 mg.  
Magnesium carbonate . . . . . 60 mg.

Binder controls and  
extends antacid activity



Egraine†‡ . . . . . 45 mg.



AVAILABLE IN BOXES OF 100 TABLETS, SPECIALLY STRIPPED FOR EASY CARRYING

\*Cyamopsis tetragonoloba gum

†Protein binder from oat

‡Trade Marks

**Organon** INC. • ORANGE, N. J.



# Preception

simple, effective conception control



PRECEPTIN<sup>®</sup>  
vaginal gel




# 3 STEPS

## using radioiodine ( $I^{131}$ )


The "MEDIAC" makes measurements of thyroid activity a simple, straightforward procedure, requires no computations—takes but a few minutes. AEC requirements can be *easily* met by any physician, and the equipment cost is lower than most doctors think. The evaluation is more accurate than the BMR test, easier than the serum protein bound iodine level, is rapidly becoming routine in hospitals, clinics and the doctor's office.

Write today for Catalog "O" with complete details on the "MEDIAC" and other medical instruments for use with radioisotopes.


Nuclear Instrument and Chemical Corporation,  
259 West Erie Street, Chicago 10, Illinois



Patient is given a radiolodine capsule or solution by mouth. An identical dose is set aside as a "standard."



After a time lapse (usually 24 hours), the "standard" is measured\* in the phantom neck. A single control then adjusts the "MEDIAC" for the patient.



Phantom is removed and activity in patient's thyroid is measured.\* Uptake of radioiodine is directly related to thyroid function and shown as a per cent of the administered dose.

\*Maximum time involved—2 minutes.

first in advantages  
for low-salt therapy

# DIASAL<sup>®</sup>

salt substitute of choice

for congestive failure  
essential hypertension  
obesity  
and whenever low salt is indicated

TASTE  
POUR-QUALITY  
APPEARANCE  
STABILITY

help keep patients on low-sodium diets

FREEDOM FROM  
UNPLEASANT  
AFTERTASTE

Keeps food attractive —  
prevents nutritional deficiencies

OPTIMAL  
POTASSIUM  
CHLORIDE  
CONTENT

avoids borderline hypopotassemia — makes DIASAL  
"...the product of choice for this purpose."\*

DIASAL contains only potassium chloride, glutamic acid and inert ingredients...no sodium, lithium, or ammonium... and it is safe for prolonged use, both at the table and in cooking.

packaging: available in 2-ounce shakers and 8-ounce bottles. Send for liberal supplies of tasting samples and low-sodium diet sheets for your patients.

\*Fremont, R. E.; Rimmerman, A. B., and Shaftel, H. E.: Postgrad. Med. 10:216, 1951.

**E. FOUGERA & CO., INC.** 75 Varick St., N. Y. 13, N. Y.

A welcome change from "No" to "Oh"



## 'Sulfa-Neolin'

(BENZATHINE PENICILLIN—G WITH SULFONAMIDES, LILLY)

*chocolate-mint-flavored  
suspension for  
the treatment of  
mixed infections  
caused by  
penicillin-susceptible  
and/or  
sulfonamide-sensitive  
organisms*

The twofold antibacterial action of 'Sulfa-Neolin' is particularly effective in mixed infections. The combined action of penicillin and the sulfonamides is also of value in combating relatively resistant organisms. Since the mode of attack of penicillin differs from that of the sulfonamides, it is more difficult for an organism to develop resistance to the combination than to either drug used alone. The tasty chocolate-mint flavor of 'Sulfa-Neolin' quickly wins youngsters' approval. They'll like it!

Each 5 cc. (approximately 1 teaspoonful) contain sulfa: diazine, merazine, methazine, 0.167 Gm. each, and benzathine penicillin—G, 300,000 units. 'Sulfa-Neolin' is stable at room temperature for two years.

Dosage: Average dose is 1 teaspoonful four times a day. Suspension 'Sulfa-Neolin' is supplied in 60-cc. bottles (M-120).

ELI LILLY AND COMPANY • INDIANAPOLIS 6, INDIANA, U.S.A.

# Searle MICTINE Provides Effective Oral, Non-Mercurial Diuresis

The result of many years of research, Mictine, brand of aminometramide, supplies a long-felt need for an improved oral diuretic. Mictine, 1-allyl-3-ethyl-6-aminotetrahydropyrimidinedione, is not a mercurial, xanthine or sulfonamide.

**Effectiveness:** Every method for measuring the diuretic effect in man now available,

is no risk of acidosis. On high dosage, Mictine causes some side effects in some patients but on three tablets daily these side effects (anorexia and nausea, rarely vomiting, diarrhea or headache) are minimal or absent.

**Indications:** Mictine is useful primarily in the *maintenance* of an edema-free state and in the *initial and continuing* control of

patients in mild congestive failure. Mictine may be used also for *initial and continuing* diuresis in *more severe* congestive states, particularly when mercurial diuretics are contraindicated.


**Administration:** The usual dosage for the average patient is one to four tablets daily with meals, in divided

doses on an interrupted schedule. An interrupted dosage schedule may be accomplished by giving the drug on alternate days or for three consecutive days and then omitting it for four days.

For severe congestive states the dosage is four to six tablets daily with meals, in divided doses on interrupted schedules similar to those already mentioned.

**Supplied:** Uncoated tablets of 200 mg.

\*Trademark of G. D. Searle & Co.




*Mictine is believed to act by the selective inhibition of the reabsorption of sodium ions. Thus, the resulting diuresis is characterized by increased quantities of sodium ions and water.*

including precise human bioassay studies, without exception demonstrated that Mictine is an effective oral diuretic, and these studies show that approximately 70 per cent of *unselected* edematous patients treated with Mictine by mouth respond with a satisfactory diuresis.

**Well-Tolerated:** There are no known contraindications to Mictine, even in the presence of hepatic or renal damage, and there

SEARLE



...through  
the perilous  
night

## You can prevent attacks in angina pectoris

Fear is a faithful companion. In angina pectoris, particularly, many patients live in constant dread of recurrent attacks.

Prophylaxis with Peritrate, a long-acting coronary vasodilator, offers new security in a majority of such cases. A single dose affords protection for as long as 4 to 5 hours, compared to 30 minutes or less with nitroglycerin.

Different investigators<sup>1-3</sup> observed that 80% of their patients responded to Peritrate therapy with fewer, less severe attacks . . . reduced nitroglycerin dependence . . . improved EKG's.

A variety of convenient dosage forms now extends these benefits. Peritrate Delayed Action tablets (10 mg.), taken with the regular bedtime dose of Peritrate (plain) help allay the fear of nighttime attacks. Adapted to the recommended daily dosage of 40-80 mg., Peritrate is available in 10 mg. and 20 mg. tablets. And when added sedation is indicated, you can prescribe Peritrate (10 mg.) with Phenobarbital (15 mg.).

1. Winsor, T., and Humphreys, P.: *Angiology* 3:1 (Feb.) 1952. 2. Plotz, M.: *New York State J. Med.* 52:2012 (Aug. 15) 1952. 3. Dailheu-Geoffroy, P.: *L'Ouest-Médical*, vol. 3 (July) 1950.

# Peritrate®

tetranitrate

(BRAND OF PENTAERYTHRITOL TETRANITRATE)

**WARNER-CHILCOTT**

# NEW! 'Tetrazets'

BACITRACIN-TYROTHRIN-NEOMYCIN-BENZOCAINE TROCHES

*broader attack to overcome minor throat irritations*

**MAJOR ADVANTAGES:** Combines 3 antibiotics to fight both gram-positive and gram-negative bacteria. Benzocaine included for soothing effect. Little danger of sensitization.



'TETRAZETS' quickly relieve minor mouth and throat irritations

*It's new*—a single troche containing 3 potent antibiotics (bacitracin, tyrothricin, neomycin) to combat afebrile oral infections.

'TETRAZETS' offer you the ideal topical treatment of minor irritations of the oral cavity.

In deep-seated infections, such as Vincent's infection, tonsillitis and streptococcus sore throat, 'TETRAZETS' may be used as an adjuvant to parenteral antibiotics.

Before and after tonsillectomies, 'TETRAZETS' help combat secondary invaders.

*Supplied:* In vials of 12. Each 'TETRAZETS' troche contains 50 units of zinc bacitracin, 1 mg. tyrothricin, 5 mg. neomycin sulfate with 5 mg. benzocaine.



Philadelphia 1, Pa.  
DIVISION OF MERCK & CO., INC.

ed—a suitable dosage form

**CAPSULES**

*Lederle*

\* REG. U. S. PAT. OFF.

**Other Dosage Forms:**

**TABLETS:** 50, 100, and 250 mg.

**PEDIATRIC DROPS** (Cherry Flavor): 100 mg. per cc. (approx. 5 mg. per drop), 10 cc. bottle

**ORAL SUSPENSION** (Cherry Flavor): 250 mg. per teaspoonful (5 cc.), 1 oz. bottle

**SPERSOIDS\*** Dispersible Powder (Chocolate Flavor): 50 mg. per rounded teaspoonful (3 Gm.), 12 and 25 dose bottle

**SOLUBLE TABLETS:** 50 mg.

**INTRAVENOUS:** vials of 100, 250, and 500 mg.

**INTRAMUSCULAR:** vial of 100 mg.

**OINTMENT** (3%):  $\frac{1}{2}$  and 1 oz. tubes

**OPHTHALMIC OINTMENT** (1%):  $\frac{1}{2}$  oz. tubes

**OPHTHALMIC SOLUTION:** vial of 25 mg. with sterilized dropper vial

**EAR SOLUTION** (0.5%): 10 cc. dropper bottle

**SYRUP** (Cherry Flavor): 125 mg. per teaspoonful (5 cc.), 2 oz. bottle

In any of its numerous dosage forms, ACHROMYCIN is a potent weapon to combat a variety of infections. This true broad-spectrum antibiotic has proved effective in controlling Gram-negative and Gram-positive bacteria, rickettsia, spirochetes, and certain viruses and protozoa.

And ACHROMYCIN is unusually kind to the patient! Extensive use, in children and adults, has shown it to be practically free of untoward side reactions. ACHROMYCIN diffuses rapidly in body tissues and fluids, is readily soluble, and remains stable for 24 hours in solution.

When a broad-spectrum antibiotic is indicated, remember—for your convenience, for the patient's comfort, there's a particular dosage form of ACHROMYCIN to fill the need!

LEDERLE LABORATORIES DIVISION  
*AMERICAN Cyanamid COMPANY* PEARL RIVER, NEW YORK

Now available as Syrup—ready-mixed!

**ACHROMYCIN\***

TETRACYCLINE LEDERLE



# Pure as sunlight



**Upjohn**

**Sex hormones—  
only one injection  
per month:**

---

## **Depo-Estradiol**

Trademark, Reg. U. S. Pat. Off.

CYCLOPENTYLPROPIONATE

*Each cc. contains:*

Estradiol, 17-Cyclopentylpropionate .....	1 mg. or 5 mg.
Chlorobutanol .....	5 mg.
Cottonseed Oil .....	q.s.
1 mg./cc. strength in 10 cc. vials	
5 mg./cc. strength in 5 cc. vials	

## **Depo-Testosterone**

Trademark, Reg. U. S. Pat. Off.

CYCLOPENTYLPROPIONATE

*Each cc. contains:*

Testosterone Cyclopentylpropionate .....	50 mg. or 100 mg.
Chlorobutanol .....	5 mg.
Cottonseed Oil .....	q.s.
50 mg./cc. strength in 10 cc. vials	
100 mg./cc. strength in 1 cc. and 10 cc. vials	

## **Depo-Testadiol**

Trademark, Reg. U. S. Pat. Off.

*Each cc. contains:*

Testosterone Cyclopentylpropionate .....	50 mg.
Estradiol, 17-Cyclopentylpropionate .....	2 mg.
Chlorobutanol .....	5 mg.
Cottonseed Oil .....	q.s.
10 cc. vials	

**The Upjohn Company, Kalamazoo, Michigan**

**"Premarin" relieves  
menopausal symptoms with  
virtually no side effects, and  
imparts a highly gratifying  
"sense of well-being."**

"Premarin"® — Conjugated Estrogens (equine)

11-51

## PLASTIC AND GLASS EYES



## MAGER & GOUGELMAN, Inc.

OVER A CENTURY OF SERVICE • 1851-1952

- PRIVATE FITTINGS
- PERSONAL ATTENTION
- EXPERIENCE & SKILL

**510 MADISON AVE., N. Y. 22**  
Telephone PLaza 5-3756

**when proper or corrective  
shoes are indicated . . .**

**prescribe PEDIFORMES**

### IN STOCK:

- straight last shoes
- inflare last shoes
- outflare last shoes
- club foot shoes
- flat foot shoes
- children's shoes with Thomas  
Heel and long counter

You are assured alterations and fittings as you prescribe because of our interest, and experience gained in serving the medical profession for more than 40 years.



34 West 36th STREET  
New York City  
Wisconsin 7-2534

Other Shops in: Brooklyn • Flatbush • Hempstead  
New Rochelle • Hackensack

# The long and short of Bentyl's relief of nervous gut

Clinicians<sup>1,2</sup> prove Bentyl is long on effective relief... short on unwanted side effects including blurred vision and dry mouth.

1. McHardy and Browne: Sou. Med. J. 45:1139, 1952. 2. Lorber and Shay: Fed. Proc. 12:90, 1952.

Complete Bentyl bibliography on request.

(Dicyclomine Hydrochloride)

another exclusive development of Merrell research

## Rx INFORMATION

### BENTYL

Bentyl affords direct (musculo-tropic) and indirect (neuro-tropic) spasmolytic action. Bentyl provides complete and comfortable relief in smooth muscle spasm; particularly in functional G.I. disorders, in irritable colon, pylorospasm, biliary tract dysfunction and spastic constipation.

**Composition:** Each capsule or teaspoonful (5 cc.) contains 10 mg. of Bentyl (dicyclomine hydrochloride).

Bentyl with Phenobarbital adds 15 mg. of phenobarbital to the preceding formula.

**Dosage:** Adults, 2 capsules or 2 teaspoonfuls of syrup, three times daily, before or after meals. If necessary, repeat dose at bedtime. In Infant Colic,  $\frac{1}{2}$  to 1 teaspoonful, ten to fifteen minutes before feeding.

**Supplied:** Bentyl — In bottles of 100 and 500 blue capsules, and Bentyl Syrup in pint and gallon bottles. Bentyl with Phenobarbital — In bottles of 100 and 500 blue-and-white capsules, and Bentyl Syrup in pint and gallon bottles.

T.M. "BENTYL"



PIONEER IN MEDICINE FOR OVER 125 YEARS

THE WM. S. MERRELL COMPANY New York • CINCINNATI • St. Thomas, Ontario

agents  
of *for the treatment of pneumonia*  
*and other respiratory tract infections*  
choice  
Terramycin<sup>®</sup>  
BRAND OF OXYTETRACYCLINE


For (ESTABLISHED) broad-spectrum antibiotic therapy—supplied in convenient Capsules, Tablets (sugar coated), Oral Suspension (raspberry flavored), Pediatric Drops (raspberry flavored), Intramuscular, Intravenous, Ophthalmics, Ointment and other topical forms.

Tetracyn<sup>®</sup>  
BRAND OF TETRACYCLINE

For the (NEWEST) broad-spectrum antibiotic therapy—supplied in convenient Capsules, Tablets (sugar coated), Oral Suspension (chocolate flavored), Pediatric Drops (banana flavored), Intramuscular, Intravenous, Ophthalmic and Ointment.

Both discovered by  world's largest producer of antibiotics

Pfizer Laboratories, Brooklyn 6, N. Y.  
Division, Chas. Pfizer & Co., Inc.



## peptic ulcer

In a recent study of the basal secretion of duodenal ulcer patients,<sup>1</sup> a single 10 mg. oral dose of Monodral was found to produce anacidity in 38 of 47 tests and to reduce the volume of secretion to a few milliliters in 33.

**Heals, does not conceal.** Monodral relieves ulcer pain faster, and favors healing, because it eliminates two pain-producing factors: excess free acid and gastric hypermotility.<sup>2</sup>

Anticholinergic agents which do not have this ability to suppress HCl may mask the failure of the ulcer to heal. Try Monodral on patients who have failed to respond to other parasympatholytic agents. Milder side effects—virtually no constipation.

**dral<sup>®</sup>** bromide

1. McKenna, R.D.: Royal Victoria Hospital, Montreal, Canada: Personal communication.

2. Ingegno, A.P.; and Kertzner, Leonard: New York Jour. Med., 54:1185, Apr. 15, 1954.

*Winthrop-Stearns* INC.

New York 18, N. Y. Windsor, Ont.

MONODRAL BROMIDE, BRAND OF PERTHLENATE BROMIDE

# CHARLES B. TOWNS HOSPITAL Established 1901

**FOR THE TREATMENT OF ALCOHOLISM**

**NARCOTIC AND BARBITURATE**

**ADDICTIONS EXCLUSIVELY**


Complete Medical & Psychiatric Treatment at predetermined cost. Privacy of patient is assured—if desired.

Literature on Request

Edward B. Towns, Director

293 Central Park West, N. Y. 24, N. Y.  
Schuyler 4-0770

Member American Hospital Association



**SATISFACTION GUARANTEED**

**Specialists in Plastic and Glass  
Artificial Human Eyes Exclusively**

*Referred Cases Carefully Attended*

**FRIED & KOHLER, Inc.**

665 FIFTH AVENUE near 53rd Street  
NEW YORK, N. Y. Tel. ELdorado 5-1970

*For acidosis due to anesthesia — edema*

## KALAK Counter-Acts ANTI-BIOTIC REACTIONS



. . . . KALAK is a non-laxative, alkaline diuretic buffer — side reactions from aureomycin — terramycin — sulfas — penicillin are reduced through the use of KALAK — KALAK contains only those salts **NORMALLY** present in plasma. . . IT IS BASIC!

**KALAK WATER CO.  
of NEW YORK, Inc.**

90 West St., New York 6, N. Y.

*For acidosis due to nausea — in nephritis*

SEDATION  
AND EUPHORIA FOR NERVOUS,  
IRRITABLE PATIENTS

Use

## VALERIANETS-DISPERT

Reg. U. S. Pat. Off.  
Each Chocolate Coated Tablet Contains Ext. Valerian (highly concentrated) 0.05 Gm. disintegrating finely subdivided for maximum efficiency  
**TASTELESS, ODORLESS, NON-DEPRESSANT SEDATIVE and EUPHORIC**  
VALERIANETS-DISPERT are indicated in cases of nervous excitement and exhaustion, anxiety and depressive states, cardiac and gastrointestinal neuroses, menopausal and menstrual melenia, insomnia.

Dose: 1 or 2 tablets t.i.d. — Bottle of 50 and of 100 tablets  
At All Prescription Pharmacies



STANDARD PHARMACEUTICAL CO., INC., 253 West 26th St., New York

For Intestinal Dysfunction  
**NUCARPON®**

Each tablet cont: Extract of Rhubarb, Senna, Precip. Sulfur, Peppermint Oil, Fennel Oil in activated charcoal base.

For making Burrow's Solution  
U.S.P. XIV  
WET DRESSING Use

**PRESTO-BORO®**

(Aluminum Sulfate and Calcium Acetate)  
POWDER IN ENVELOPES  
— TABLETS —

For treatment of Swellings, Inflammations, Sprains

For Pulmonary Conditions  
**TRANSPULMIN®**

3% solution Quinine with 2 1/2% Camphor for intramuscular injection.

# BETTER TOLERATED SALICYLATE THERAPY

**For Headache, Neuralgia, Minor Aches and Pains**  
**Give BUFFERIN® because...**

1. It gives fast pain relief—acts twice as fast as aspirin.<sup>1</sup>
2. Even large doses seldom cause gastric upsets.<sup>2</sup>

**For Arthritis—and Other Rheumatic Disorders**  
**Give BUFFERIN because...**

1. It provides effective, better-tolerated relief of pain.
2. There were no gastric upsets with BUFFERIN in 70% of hospitalized arthritic patients who couldn't tolerate aspirin.<sup>3</sup> This is an important finding, for arthritics are 3 to 9 times as susceptible to gastric upsets with straight aspirin as the general population.<sup>3</sup>
3. The antacids in BUFFERIN do not lower the blood salicylate levels as sodium bicarbonate does.<sup>4</sup>



BUFFERIN contains acetylsalicylic acid (5 gr. per tablet), for prompt analgesia, plus magnesium carbonate and aluminum glycinate.

**Available**—bottles of 12, 36, 60 and 100 tablets.

**References:** 1. J. Am. Pharm. Assoc., Sc. Ed. 39:21 (Jan.) 1950. 2. Ind. Med. 20:480 (Oct.) 1951. 3. In Press. 4. J.A.M.A. 141:124 (Sept. 10) 1949.

**WHENEVER SALICYLATE THERAPY IS INDICATED**  
**GIVE BUFFERIN® Acts Twice as Fast as Aspirin**  
**Does Not Upset the Stomach**

**BRISTOL-MYERS CO.,**  
**19 W. 50 St., New York 20, N. Y.**



**a pain-free stomach—  
quiet intestines—  
for your nervous patients**

**'TRICOLOID'\***

quickly blocks the nervous  
impulses which cause gastrointestinal  
symptoms;

causes negligible dryness of the mouth or  
other side effect.

**\*'TRICOLOID' brand Tricyclamol 50 mg. Compressed, sugar-coated  
Bottles of 100**



**BURROUGHS WELLCOME & CO. (U.S.A.) INC., Tuckahoe 7, New York**

**broad spectrum antibiotic of choice**

# STECLIN

SQUIBB TETRACYCLINE

**reaches higher levels in the urine than its analogues**

**crosses the intact meningeal  
barrier more readily than its analogues**

**produces higher blood levels than its analogues**

**fewer gastrointestinal side effects than its analogues**

***Minimum adult dose: 250 mg. q.i.d.***

**Capsules: 250 mg., bottles of 16 and 100.  
50 and 100 mg., bottles of 25 and 100.**

**Suspension: 1 oz. bottles, supplied with dropper calibrated at 1 cc.  
Each 5 cc. teaspoonful contains the equivalent of 250 mg.  
of tetracycline hydrochloride.**



\*STECLIN\* IS A SQUIBB TRADEMARK

**Hoeber-Harper**



**announces . . .**

## **THREE IMPORTANT *new* BOOKS**

**BAKER'S:**

---

### ***Clinical Neurology***

---

*In 3 Volumes.* A complete source of today's knowledge of diagnosis and treatment—including not only the more common neurologic disturbances but also the rarer syndromes and the ramifications of neurology in the fields of internal medicine, pediatrics, endocrinology, psychiatry, ophthalmology and otolaryngology. While the first objective is to organize for practical use the known facts of neurology, much new material is published here for the first time.

*65 Authorities. Edited by A. B. BAKER, M.D., Prof. and Dir., Div. of Neurology, Univ. of Minn. Med. Sch., 2100 pp., liberally illustrated, 3-volumes \$60.00.*

**WERNER'S:**

---

### ***The Thyroid***

---

This interdisciplinary approach to the thyroid gland provides the first complete guide to thyroid functions, disorders and relationships. From the fundamental biochemistry to the details of treatment, both medical and surgical, including laboratory technics as well as the latest radioactive materials, you will find today's best knowledge on any question you may look up. Each topic is presented by an authority; yet, because the book is so carefully planned and edited, it reads like a consecutive account by a single authority.

*53 Authors. Edited by SIDNEY C. WERNER, M.D., Assoc. Prof. of Clinical Medicine, Columbia Univ. Coll. of Physicians and Surgeons. Approx. 800 pp. In Press.*

**DUNHAM'S:**

---

### ***Premature Infants***

---

**2nd Edition**

*A Manual for Physicians.* The basic source of information on prematurity and guide to care of the premature infant. Part I deals with general considerations: criteria for prematurity; incidence, causes and prevention of premature birth; growth and development of premature infants. Part II takes up the physiologic handicaps of premature infants, their problems and their care; congenital and acquired conditions that tend to affect them adversely. Part III outlines public health aspects of prematurity, including death rates and causes of death.

*ETHEL C. DUNHAM, M.D., Children's Bureau. In Press.*

**All Hoeber-Harper Books Sent On Approval**

---

**PAUL B. HOEBER, Inc., Publisher, 49 East 33rd St., N.Y. 16**

**MEDICAL BOOK DEPARTMENT OF HARPER & BROTHERS**